

A NOTE ON REGULARITY OF SOLUTIONS TO DEGENERATE ELLIPTIC EQUATIONS OF CAFFARELLI-KOHN-NIRENBERG TYPE

VERONICA FELLI AND MATTHIAS SCHNEIDER

ABSTRACT. We establish Hölder continuity of weak solutions to degenerate critical elliptic equations of Caffarelli-Kohn-Nirenberg type.

1. INTRODUCTION

Our purpose is to establish Hölder continuity of weak solutions to

$$-\operatorname{div}(|x|^{-2a}\nabla u) = \frac{f}{|x|^{bp}}, \text{ in } \Omega \subset \mathbb{R}^N, \quad (1.1)$$

where Ω is an open, $N \geq 3$ and a , b , and p satisfy

$$\begin{aligned} -\infty < a < \frac{N-2}{2}, \quad a \leq b \leq a+1 \\ p = p(a, b) = \frac{2N}{N-2(1+a-b)}. \end{aligned} \quad (1.2)$$

We denote by $\mathcal{D}_a^{1,2}(\Omega)$ the closure of $C_c^\infty(\Omega)$ with respect to the norm

$$\|u\|_{\mathcal{D}_a^{1,2}(\Omega)} := \left(\int_{\Omega} |\nabla u|^2 |x|^{-2a} \right)^{1/2}.$$

For a given weight ω we denote by $L^p(\Omega, \omega)$ the space of functions u such that

$$\|u\|_{L^p(\Omega, \omega)}^p := \int_{\Omega} |u|^p \omega(x) < \infty.$$

The space $H_a^1(\Omega)$ is defined to be the closure of $C^\infty(\bar{\Omega})$ with respect to

$$\|u\|_{H_a^1(\Omega)}^2 := \int_{\Omega} |x|^{-2a} (|\nabla u|^2 + |u|^2).$$

Our interest in these problems arose because of their relation to nonlinear, degenerate elliptic equations stemming from the family of Caffarelli-Kohn-Nirenberg inequalities [2]: if a , b , and p satisfy (1.2) then we have for all $u \in \mathcal{D}_a^{1,2}(\mathbb{R}^N)$

$$\left(\int |u|^p |x|^{-bp} \right)^{1/p} \leq C_{a,b,N} \left(\int |\nabla u|^2 |x|^{-2a} \right)^{1/2}. \quad (1.3)$$

Date: June 25, 2003.

1991 *Mathematics Subject Classification.* 35J70, 35B65, 35B45.

Key words and phrases. critical exponents, Caffarelli-Kohn-Nirenberg inequality, degenerate elliptic.

V. F. is supported by M.U.R.S.T. under the national project “Variational Methods and Nonlinear Differential Equations”. During the preparation of the paper M. S. research was supported by a S.I.S.S.A. postdoctoral fellowship.

For best constants and existence of minimizers in (1.3) we refer to [3]. Due to its characterization any minimizer $u \in \mathcal{D}_a^{1,2}(\mathbb{R}^N)$, if it exists, is a weak solution to

$$-\operatorname{div}(|x|^{-2a}\nabla u) = \frac{K(x)|u|^{p-2}u}{|x|^{bp}} \text{ in } \mathbb{R}^N, \quad (1.4)$$

where $K(x) \equiv \text{const}$ for some appropriate constant. The exponent $p = p(a, b, N)$ is the critical exponent in (1.3) and shares many features with the critical Sobolev exponent, e.g. (1.4) possesses for $K(x) \equiv K(0)$ a dilation symmetry, which gives rise to a noncompact manifold of weak radial solutions for $K(0) > 0$. In order to study problem (1.4) for non-constant functions K using for instance a degree argument, Hölder estimates for weak solutions of (1.1) are an important tool (see [7]).

Regularity properties of weak solution to degenerate elliptic problems with more general weighted operators of the form $\operatorname{div}(\omega(x)\nabla(\cdot))$ are studied in [5, 6, 8] (see also the references mentioned there). The classes of weights ω treated there include the class (QC') of weights

$$\omega(x) = |\det T'|^{1-2/N},$$

where $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is quasi-conformal (see [6, 8] for a definition). In fact our weights $|\cdot|^{-2a}$ are associated with quasi-conformal transformations $T_a(x) := x|x|^{-2a/(N-2)}$. The right-hand sides studied in [5, 6, 8] are either zero or in divergence form, e.g. Hölder continuity of weak solutions to

$$-\operatorname{div}(|x|^{-2a}\nabla u) = \operatorname{div}(F) \text{ in } \Omega$$

is established in [5] assuming $|F||x|^{2a} \in L^p(\Omega, |x|^{-2a})$ for some $p > \max(N - 2a, N, 2)$. We derive Hölder estimates for weak solutions to (1.1) in terms of f , because a sharp relation of the integrability of f and its representation in divergence form F in the various weighted spaces is not obvious. We compare weak solutions of (1.1) with μ_a -harmonic functions, which are by definition weak solutions of

$$-\operatorname{div}(|x|^{-2a}\nabla u) = 0 \text{ in } \Omega,$$

where Hölder regularity is known (see for instance [8]) and prove

Theorem 1.1. *Suppose $\Omega \subset \mathbb{R}^N$ is a bounded domain and $u \in H_a^1(\Omega)$ weakly solves (1.1), that is*

$$\int_{\Omega} |x|^{-2a} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} |x|^{-bp} f \varphi \, dx \quad \forall \varphi \in H_{0,a}^1(\Omega).$$

Assume a, b and p satisfy (1.2), $b < a + 1$, and $f \in L^s(\Omega, |x|^{-bp})$ for some $s > p/(p - 2)$. Then $u \in C^{0,\alpha}$ for any $\alpha \in (0, 1)$ satisfying

$$\alpha < \min(\alpha_h, 1) \text{ and } \alpha < \begin{cases} (\frac{N-2}{2} - a)(p - 2 - \frac{p}{s}) & \text{if } b \geq 0 \\ \frac{N}{p}(p - 2 - \frac{p}{s}) & \text{if } b < 0 \end{cases},$$

where α_h is the regularity of μ_a -harmonic functions given in Theorem 2.1 below. Moreover, for any $\Omega' \Subset \Omega$ there is a constant $C = C(N, a, \alpha, \Omega, \operatorname{dist}(\Omega', \Omega), s)$ such that

$$\sup_{\Omega'} |u| + \sup_{\substack{x, y \in \Omega' \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq C \left\{ \|u\|_{L^2(\Omega, d\mu_a)} + \|f\|_{L^s(\Omega, |x|^{-bp})} \right\}$$

For the nonlinear problem (1.4) we use a De Giorgi-Moser type iteration procedure as in [1] and obtain

Theorem 1.2. *Let a, b and p satisfy (1.2) and $u \in D_a^{1,2}(\mathbb{R}^N)$ be a weak solution to*

$$-\operatorname{div}(|x|^{-2a}\nabla u) = K(x)\frac{|u|^{p-2}u}{|x|^{bp}}, \quad x \in \Omega \quad (1.5)$$

where $K \in L^\infty(\Omega)$. Then $u \in L_{\text{loc}}^s(\Omega, |x|^{-bp})$ for any $s \in [p, +\infty[$. Moreover, u is Hölder continuous with Hölder exponent given in Theorem 1.1.

Remark 1.3. While completing this note we learned that in [4] weighted q -Laplacian equations of the form

$$-\operatorname{div}(|x|^{-qa}|\nabla u|^{q-2}\nabla u) = g$$

are studied. Under assumption (1.2) Hölder regularity of weak solutions to equation

$$-\operatorname{div}(|x|^{-qa}\nabla u) = \frac{f}{|x|^{bp}}, \quad \text{in } \Omega \subset \mathbb{R}^N,$$

is shown if $a = b$, $a > -1$, and $f \in L^s(\Omega, |x|^{-bp})$ for some $s > p/(p-2)$. Theorem 1.1 extends this result to the full range for a and b in the case $q = 2$.

2. PRELIMINARIES

We collect some properties of the weighted measure $\mu_a := |x|^{-2a} dx$ and μ_a -harmonic functions. We refer to [6, 8] for the proofs.

- The measure μ_a satisfies the doubling property, i.e. for every $\tau \in (0, 1)$ there exists a constant $C_{(2.1)}(\tau)$ such that

$$\mu_a(B(x, r)) \leq C_{(2.1)}(\tau)\mu_a(B(x, \tau r)) \quad (2.1)$$

- A Poincaré-type inequality holds, i.e. there is a positive constant $C_{(2.2)}$ such that any $u \in D_a^{1,2}(\mathbb{R}^N)$ satisfies

$$\int_{B_r(x)} |u - u_{x,r}|^2 d\mu_a \leq C_{(2.2)} r^2 \int_{B_r(x)} |\nabla u|^2 d\mu_a, \quad (2.2)$$

where $u_{x,r}$ denotes the weighted mean-value

$$u_{x,r} := \int_{B_r(x)} u d\mu_a = \frac{1}{\mu_a(B_r(x))} \int_{B_r(x)} u(x) d\mu_a.$$

Concerning μ_a -harmonic functions we have the following results.

Theorem 2.1 (Thm. 3.34 in [8](p. 65), Thm. 6.6 in [8](p. 111)).

There are constants $C_{(2.3)}(N, a)$ and $\alpha_h = \alpha_h(N, a) \in (0, 1)$ such that if u is μ_a -harmonic in $B_r(x_0) \subset \mathbb{R}^N$ and $0 < \rho < r$ then

$$\operatorname{ess-sup}_{B(x_0, \frac{r}{2})} |u| \leq C_{(2.3)} \int_{B(x_0, r)} |u|^2 d\mu_a, \quad (2.3)$$

$$\operatorname{osc}(u, B_\rho(x_0)) \leq 2^{\alpha_h} \left(\frac{\rho}{r}\right)^\alpha \operatorname{osc}(u, B_r(x_0)). \quad (2.4)$$

Consequently, μ_a -harmonic functions are Hölder continuous.

We will call a function $u \in D_{a, \text{loc}}^{1,2}(\mathbb{R}^N)$ weakly super μ_a -harmonic in Ω , if for all nonnegative $\varphi \in C_c^\infty(\Omega)$ we have

$$\int_{\Omega} |x|^{-2a} \nabla u \nabla \varphi \geq 0. \quad (2.5)$$

Theorem 2.2 (Thm 3.51 in [8](p. 70)). *There exist positive constants $s = s(N, a)$ and $C_{(2.6)} = C_{(2.6)}(N, a)$ such that if u is nonnegative and weakly super μ_a -harmonic in Ω and $B_{2r}(x_0) \subset \Omega$ we have*

$$\operatorname{ess\,inf}_{B_{\frac{r}{2}}(x_0)} u \geq C_{(2.6)} \left(\int_{B_r(x_0)} u^s d\mu_a \right)^{\frac{1}{s}}. \quad (2.6)$$

We use the two theorems above to derive

Lemma 2.3. *For any ball $B_r(x_0)$ there is a constant $C_{(2.7)}(B_r(x_0))$ such that any μ_a -harmonic function u in $B_r(x_0)$ satisfies*

$$\int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a \leq C_{(2.7)} \left(\frac{\rho}{r} \right)^{2\alpha_h - 2} \int_{B_r(x_0)} |\nabla u|^2 d\mu_a, \quad (2.7)$$

where $\alpha_h \in (0, 1)$ is given in Theorem 2.1.

Proof. To prove the claim we may assume that $0 < \rho < (1/4)r$ and that u has mean-value zero in $B_r(x_0)$. We take a cut-off function $\xi \in C_c^\infty(B_{2\rho}(x_0))$ such that $\xi \equiv 1$ in $B_\rho(x_0)$, $0 \leq \xi \leq 1$, $\|\nabla \xi\|_\infty \leq 2\rho^{-1}$ and define $\varphi := \xi^2(u - u(x_0))$. Testing with φ and using Hölder's inequality we get

$$\int_{B_r(x_0)} |\nabla u|^2 \xi^2 d\mu_a \leq \int_{B_r(x_0)} |\nabla \xi|^2 (u - u(x_0))^2 d\mu_a \leq \|u - u(x_0)\|_{\infty, B_{2\rho}(x_0)}^2 \mu_a(B_{2\rho}(x_0)) \rho^{-2}. \quad (2.8)$$

From (2.8) and (2.4) we infer

$$\int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a \leq C \left(\frac{\rho}{r} \right)^{2\alpha} \operatorname{osc}(u, B_{\frac{r}{2}}(x_0))^2 \rho^{-2} \leq C \left(\frac{\rho}{r} \right)^{2\alpha} \rho^{-2} \int_{B_r(x_0)} |u|^2 d\mu_a$$

Finally, since u has mean-value zero in $B_r(x_0)$ the Poincaré inequality (2.2) yields the claim. \square

3. GROWTH OF LOCAL INTEGRALS

We give a weighted version of the Campanato-Morrey characterization of Hölder continuous functions.

Theorem 3.1. *Suppose $\Omega \subset \mathbb{R}^N$ is a bounded domain and $u \in L^2(\Omega, d\mu_a)$ satisfies*

$$\int_{B_r(x)} |u(y) - u_{x,r}|^2 d\mu_a \leq M^2 r^{2\alpha} \quad \text{for any } B_r(x) \subset \Omega \quad (3.1)$$

and some $\alpha \in (0, 1)$. Then $u \in C^{0,\alpha}(\Omega)$ and for any $\Omega' \Subset \Omega$ there holds

$$\sup_{\Omega'} |u| + \sup_{\substack{x, y \in \Omega' \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq C \left\{ M + \|u\|_{L^2(\Omega, |x|^{-2a})} \right\}$$

where $C = C(N, a, \alpha, \Omega, \operatorname{dist}(\Omega', \Omega))$ is a positive constant independent of u .

Proof. Denote $R_0 = \operatorname{dist}(\Omega', \partial\Omega)$. Using the triangle inequality and integrating in $B_{r_1}(x_0)$ we have for any $x_0 \in \Omega'$ and $0 < r_1 < r_2 \leq R_0$

$$\begin{aligned} & |u_{x_0, r_1} - u_{x_0, r_2}|^2 \\ & \leq \frac{2}{\mu_a(B_{r_1}(x_0))} \left\{ \int_{B_{r_1}(x_0)} |u(x) - u_{x_0, r_1}|^2 d\mu_a + \int_{B_{r_2}(x_0)} |u(x) - u_{x_0, r_2}|^2 d\mu_a \right\}. \end{aligned}$$

Using assumption (3.1) we obtain

$$|u_{x_0, r_1} - u_{x_0, r_2}|^2 \leq \frac{2M^2}{\mu_a(B_{r_1}(x_0))} \{ \mu_a(B_{r_1}(x_0)) r_1^{2\alpha} + \mu_a(B_{r_2}(x_0)) r_2^{2\alpha} \}. \quad (3.2)$$

For any $R \leq R_0$ we take $r_1 = 2^{-(i+1)}R$ and $r_2 = 2^{-i}R$ in (3.2). The doubling property (2.1) then gives

$$|u_{x_0, 2^{-(i+1)}R} - u_{x_0, 2^{-i}R}| \leq 2M^2 \left(1 + C_{(2.1)}(N, a) 2^{2\alpha}\right) 2^{-2(i+1)\alpha} R^{2\alpha}.$$

We sum up and get for $h < k$

$$|u_{x_0, 2^{-h}R} - u_{x_0, 2^{-k}R}| \leq \frac{C(N, a, \alpha)M}{2^{h\alpha}} R^\alpha. \quad (3.3)$$

The above estimates prove that $\{u_{x_0, 2^{-i}R}\}_{i \in \mathbb{N}} \subset \mathbb{R}$ is a Cauchy sequence in \mathbb{R} , hence it converges to some limit, denoted as $\hat{u}(x_0)$. The value of $\hat{u}(x_0)$ is independent of R , which may be seen by analogous estimates. Consequently, from (3.3) we have that

$$|u_{x_0, r} - \hat{u}(x_0)| \leq C(N, a, \alpha) M r^\alpha \quad \forall x_0 \in \Omega'. \quad (3.4)$$

By the Lebesgue theorem we infer

$$u_{x, r} = \frac{|B_r(x)|}{\int_{B_r(x)} |y|^{-2a} dy} \cdot \frac{\int_{B_r(x)} |y|^{-2a} u(y) dy}{\frac{1}{|B_r(x)|}} \xrightarrow{r \rightarrow 0^+} u(x), \quad \text{a. e. in } \Omega'.$$

Hence $\hat{u} = u$ a. e. in Ω' and (3.4) gives

$$|u_{x_0, r} - u(x_0)| \leq C(N, a, \alpha) M r^\alpha \quad \forall x_0 \in \Omega', \quad (3.5)$$

which implies that $u_{x, r}$ converges to u uniformly in Ω' . Since $x \mapsto u_{x, r}$ is a continuous function, we conclude that u is continuous in Ω' . From (3.5) we have

$$|u(x)| \leq C(N, a, \alpha) M R^\alpha + |u_{x, R}| \quad \forall x \in \Omega', \quad \forall R \leq R_0.$$

Thus u is bounded in Ω' with the estimate

$$\sup_{\Omega'} |u| \leq c(N, a, \alpha, \Omega, \text{dist}(\Omega', \Omega)) \left\{ M + \|u\|_{L^2(\Omega, |x|^{-2a})} \right\}. \quad (3.6)$$

Let us now prove that u is Hölder continuous. Let $x, y \in \Omega'$ with $|x - y| = R < \frac{R_0}{2}$. Assume that $|x| < |y|$. Then we have

$$|u(x) - u(y)| \leq |u(x) - u_{x, 2R}| + |u(y) - u_{y, 2R}| + |u_{x, 2R} - u_{y, 2R}|.$$

The first two terms are estimated by (3.5), whereas for the last term we have

$$|u_{x, 2R} - u_{y, 2R}|^2 \leq 2 \{ |u_{x, 2R} - u(\xi)|^2 + |u(\xi) - u_{y, 2R}|^2 \}$$

and integrating with respect to ξ over $B_{2R}(x) \cap B_{2R}(y) \supseteq B_R(x)$ we obtain

$$|u_{x, 2R} - u_{y, 2R}|^2 \leq \frac{2}{\mu_a(B_R(x))} \left(M^2 \mu_a(B_{2R}(x)) 2^{2\alpha} R^{2\alpha} + M^2 \mu_a(B_{2R}(y)) 2^{2\alpha} R^{2\alpha} \right).$$

Since x is closer to 0 than y , we have that $\mu_a(B_{2R}(y)) \leq \mu_a(B_{2R}(x))$ and hence

$$|u(x) - u(y)| \leq C(N, a, \alpha) M |x - y|^\alpha.$$

If $|x - y| > \frac{R_0}{2}$ we can use estimate (3.6) thus finding

$$|u(x) - u(y)| \leq 2 \sup_{\Omega'} |u| \leq c 2^\alpha \left[M + \frac{1}{R_0^\alpha} \|u\|_{L^2(\Omega, |x|^{-2a})} \right] |x - y|^\alpha.$$

The proof is thereby complete. \square

Corollary 3.2. *Suppose $\Omega \subset \mathbb{R}^N$ is a bounded domain and $u \in H_a^1(\Omega)$ satisfies*

$$\int_{B_r(x)} |\nabla u|^2 d\mu_a \leq M^2 r^{2\alpha-2} \quad \text{for any } B_r(x) \subset \Omega$$

and some $\alpha \in (0, 1)$. Then $u \in C^{0,\alpha}(\Omega)$ and for any $\Omega' \Subset \Omega$ there holds

$$\sup_{\Omega'} |u| + \sup_{\substack{x,y \in \Omega' \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq c \left\{ M + \|u\|_{L^2(\Omega, |x|^{-2a})} \right\}$$

where $c = c(N, a, \alpha, \Omega, \text{dist}(\Omega', \Omega)) > 0$.

Proof. The proof follows from Theorem 3.1 and the Poincaré type inequality in (2.2). \square

Proof of Theorem 1.1. Let $w \in u + H_{0,a}^1(B_r(x_0))$ be the unique solution to the Dirichlet problem

$$\begin{cases} -\operatorname{div}(|x|^{-2a} \nabla w) = 0 & \text{in } B_r(x_0) \\ w|_{\partial B_r(x_0)} = u. \end{cases} \quad (3.7)$$

Clearly the function $v = u - w \in H_{0,a}^1(B_r(x_0))$ weakly solves

$$-\operatorname{div}(|x|^{-2a} \nabla v) = \frac{f}{|x|^{bp}} \quad \text{in } B_r(x_0).$$

Testing the above equation with v and using Hölder's inequality and (1.3), we get

$$\int_{B_r(x_0)} |\nabla v|^2 d\mu_a \leq C_{a,b,N} \left(\int_{B_r(x_0)} |x|^{-bp} |f|^{\frac{p}{p-1}} \right)^{\frac{p-1}{p}} \left(\int_{B_r(x_0)} |\nabla v|^2 d\mu_a \right)^{\frac{1}{2}}.$$

Since $f \in L^s(\Omega, |x|^{-bp})$ for some $s > p/(p-2)$ we may use Hölder's inequality with conjugate exponents $s(p-1)/p$ and

$$\frac{s(p-1)}{s(p-1)-p} = \frac{p-1}{1+(p-2-\frac{p}{s})}$$

and Lemma A.1 with $\varepsilon = 2(p-2-p/s)/p$ to obtain

$$\begin{aligned} \int_{B_r(x_0)} |\nabla v|^2 d\mu_a &\leq C_{a,b,N}^2 \left(\int_{B_r(x_0)} |x|^{-bp} |f|^s \right)^{\frac{2}{s}} \left(\int_{B_r(x_0)} |x|^{-bp} \right)^{\frac{2}{p} + \varepsilon} \\ &\leq C r^{-2+N\varepsilon} \max(r, |x_0|)^{-bp\varepsilon} \mu_a(B_r(x_0)) \left(\int_{B_r(x_0)} |x|^{-bp} |f|^s \right)^{\frac{2}{s}}. \end{aligned} \quad (3.8)$$

From (2.7) we deduce for any $0 < \rho \leq r$

$$\begin{aligned} \int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a &\leq 4 \int_{B_\rho(x_0)} |\nabla w|^2 d\mu_a + 4 \int_{B_\rho(x_0)} |\nabla v|^2 d\mu_a \\ &\leq 4C_{(2.7)} \left(\frac{\rho}{r}\right)^{2\alpha_h-2} \int_{B_r(x_0)} |\nabla w|^2 d\mu_a + 4\mu_a(B_\rho(x_0))^{-1} \int_{B_r(x_0)} |\nabla v|^2 d\mu_a \end{aligned} \quad (3.9)$$

Since w minimizes the Dirichlet integral we may replace w in (3.9) by u . If we further estimate the integral containing v in (3.9) using (3.8) we get

$$\begin{aligned} \int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a &\leq C \left(\mu_a(B_\rho(x_0)) \mu_a(B_r(x_0))^{-1} \left(\frac{\rho}{r} \right)^{-2+2\alpha_h} \int_{B_r(x_0)} |\nabla u|^2 d\mu_a \right. \\ &\quad \left. + r^{-2+N\varepsilon} \max(r, |x_0|)^{-bp\varepsilon} \mu_a(B_r(x_0)) \|f\|_{L^s(B_r(x_0), |x|^{-bp})}^2 \right). \end{aligned}$$

We estimate the term $\max(r, |x_0|)^{-bp\varepsilon}$ by $r^{-bp\varepsilon}$ if $b \geq 0$ and in the case $b < 0$ by a constant $C(\Omega)$. For the rest of the proof we will consider the more interesting situation $b \geq 0$. The case $b < 0$ may be treated analogously.

Lemma A.2 with $\Phi(\rho) := \int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a$ gives for $0 < \rho < r \leq r_0 := \text{dist}(x_0, \partial\Omega)$

$$\begin{aligned} \int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a &\leq C(\alpha) \left(\frac{\mu_a(B_\rho(x_0))}{\mu_a(B_r(x_0))} \left(\frac{\rho}{r} \right)^{-2+2\alpha} \int_{B_r(x_0)} |\nabla u|^2 d\mu_a \right. \\ &\quad \left. + \rho^{-2+(N-bp)\varepsilon} \mu_a(B_\rho(x_0)) \|f\|_{L^s(\Omega, |x|^{-bp})}^2 \right). \end{aligned}$$

We take a cut-off function $\xi \in C_c^\infty(B_r(x_0))$ such that $\xi \equiv 1$ in $B_{r/2}(x_0)$, $0 \leq \xi \leq 1$, $\|\nabla \xi\|_\infty \leq 2r^{-1}$ and define $\varphi := \xi^2 u$. Testing with φ and using (1.3) and Hölder's inequality we get

$$\begin{aligned} \int_{B_r(x_0)} |\nabla u|^2 \xi^2 d\mu_a &\leq C_{a,b,N} \|f\|_{L^{\frac{p}{p-1}}(\Omega, |x|^{-bp})} \|\xi^2 u\|_{\mathcal{D}_a^{1,2}(\Omega)} \\ &\quad + \|u \nabla \xi\|_{L^2(B_r(x_0), d\mu_a)} \|\nabla u \xi\|_{L^2(B_r(x_0), d\mu_a)}. \end{aligned}$$

We divide by $\|\nabla u \xi\|_{L^2(B_r(x_0), d\mu_a)}$ and obtain

$$\begin{aligned} \int_{B_r(x_0)} |\nabla u|^2 \xi^2 d\mu_a &\leq \frac{2C_{a,b,N}^2 \|f\|_{L^{\frac{p}{p-1}}(\Omega, |x|^{-bp})}^2 \|\xi^2 u\|_{\mathcal{D}_a^{1,2}(\Omega)}^2}{\|\nabla u \xi\|_{L^2(B_r(x_0), d\mu_a)}^2} + \|u \nabla \xi\|_{L^2(B_r(x_0), d\mu_a)}^2 \\ &\leq \frac{2C_{a,b,N}^2 \|f\|_{L^{\frac{p}{p-1}}(\Omega, |x|^{-bp})}^2 \|\xi^2 u\|_{\mathcal{D}_a^{1,2}(\Omega)}^2}{\|\nabla u \xi\|_{L^2(B_r(x_0), d\mu_a)}^2} + 4r^{-2} \|u\|_{L^2(B_r(x_0), d\mu_a)}^2 \\ &\leq 2C_{a,b,N}^2 \|f\|_{L^{\frac{p}{p-1}}(\Omega, |x|^{-bp})}^2 \left(\frac{8r^{-2} \|u\|_{L^2(B_r(x_0), d\mu_a)}^2}{\|\nabla u \xi\|_{L^2(B_r(x_0), d\mu_a)}^2} + 2 \right) \\ &\quad + 4r^{-2} \|u\|_{L^2(B_r(x_0), d\mu_a)}^2. \end{aligned}$$

Thus

$$\int_{B_r(x_0)} |\nabla u|^2 \xi^2 d\mu_a \leq 8C_{a,b,N}^2 \|f\|_{L^{\frac{p}{p-1}}(\Omega, |x|^{-bp})}^2 + 4r^{-2} \|u\|_{L^2(B_r(x_0), d\mu_a)}^2.$$

Taking $r = r_0$ we have for $0 < \rho \leq r_0/2$

$$\int_{B_\rho(x_0)} |\nabla u|^2 d\mu_a \leq C(N, a, \Omega, r_0) \left(\int_\Omega |u|^2 d\mu_a + \|f\|_{L^{\frac{2p}{p-2}}(\Omega, |x|^{-bp} dx)}^2 \right) \rho^{-2+2\alpha}$$

From the above estimate, Corollary 3.2 and, the fact that $(N - bp)2/p = N - 2 - 2a$ we derive the desired conclusion. \square

4. A BREZIS-KATO TYPE LEMMA

As in [1] we prove the following lemma to start an iteration procedure.

Lemma 4.1. *Let $\Omega \subset \mathbb{R}^N$ be open, a , b and p satisfy (1.2), and $q > 2$. Suppose $u \in D_a^{1,2}(\mathbb{R}^N) \cap L^q(\Omega, |x|^{-bp})$ is a weak solution of*

$$-\operatorname{div}(|x|^{-2a}\nabla u) - \frac{V(x)}{|x|^{bp}}u = \frac{f(x)}{|x|^{bp}} \quad \text{in } \Omega, \quad (4.1)$$

where $f \in L^q(\Omega, |x|^{-bp})$ and V satisfies for some $\ell > 0$

$$\int_{|V(x)| \geq \ell} |x|^{-bp}|V|^{\frac{p}{p-2}} + \int_{\Omega \setminus B_\ell(0)} |x|^{-bp}|V|^{\frac{p}{p-2}} \leq \min \left\{ \frac{1}{8}C_{a,b}^{-1}, \frac{2}{q+4}C_{a,b,N}^{-1} \right\}^{\frac{p}{p-2}}. \quad (4.2)$$

Then for any $\Omega' \Subset \Omega$

$$\|u\|_{L^{\frac{pq}{2}}(\Omega', |x|^{-bp})} \leq C(\ell, q, \Omega') \|u\|_{L^q(\Omega, |x|^{-bp})} + \|f\|_{L^q(\Omega, |x|^{-bp})}. \quad (4.3)$$

If, moreover, $u \in \mathcal{D}_a^{1,2}(\Omega)$ then (4.3) remains true for $\Omega' = \Omega$.

Proof. Hölder's inequality, (1.3) and (4.2) give for any $w \in \mathcal{D}_a^{1,2}(\Omega)$

$$\begin{aligned} \int_{\Omega} |x|^{-bp}|V(x)|w^2 &\leq \ell \int_{\substack{|V(x)| \leq \ell \text{ and} \\ x \in \Omega \cap B_\ell(0)}} |x|^{-bp}w^2 + \int_{\substack{|V(x)| \geq \ell \text{ or} \\ x \in \Omega \setminus B_\ell(0)}} |x|^{-b(p-2)}|V||x|^{-2b}w^2 \\ &\leq \ell \int_{\Omega \cap B_\ell(0)} |x|^{-bp}w^2 + \left(\int_{\Omega} \frac{w^p}{|x|^{bp}} \right)^{\frac{2}{p}} \left(\int_{\substack{|V(x)| \geq \ell \text{ or} \\ x \in \Omega \setminus B_\ell(0)}} |x|^{-bp}|V|^{\frac{p}{p-2}} \right)^{\frac{p-2}{p}} \\ &\leq \ell \int_{\Omega \cap B_\ell(0)} |x|^{-bp}w^2 + \min \left(\frac{1}{8}, \frac{2}{q+4} \right) \int_{\Omega} |x|^{-2a}|\nabla w|^2. \end{aligned} \quad (4.4)$$

Suppose now that $u \in L^q(\Omega, |x|^{-bp})$. Fix $\Omega' \Subset \Omega$ and a nonnegative cut-off function η , such that $\operatorname{supp}(\eta) \Subset \Omega$ and $\eta \equiv 1$ on Ω' . Set $u^n := \min(n, |u|) \in D_a^{1,2}(\mathbb{R}^N)$ and test (4.1) with $u(u^n)^{q-2}\eta^2 \in \mathcal{D}_a^{1,2}(\Omega)$. This leads to

$$\begin{aligned} (q-2) \int_{\Omega} |x|^{-2a}\eta^2|\nabla u^n|^2(u^n)^{q-2} &+ \int_{\Omega} |x|^{-2a}\eta^2(u^n)^{q-2}|\nabla u|^2 \\ &= \int_{\Omega} |x|^{-bp}V(x)\eta^2u^2(u^n)^{q-2} + \int_{\Omega} |x|^{-bp}f\eta^2(u^n)^{q-2}u - 2 \int_{\Omega} |x|^{-2a}\nabla u\eta(u_n)^{q-2}\nabla\eta u. \end{aligned}$$

We use the elementary inequality $2ab \leq 1/2a^2 + 4b^2$ and obtain

$$\begin{aligned} (q-2) \int_{\Omega} |x|^{-2a}\eta^2|\nabla u^n|^2(u^n)^{q-2} &+ \frac{1}{2} \int_{\Omega} |x|^{-2a}\eta^2(u^n)^{q-2}|\nabla u|^2 \\ &\leq \int_{\Omega} |x|^{-bp}V(x)\eta^2u^2(u^n)^{q-2} + \int_{\Omega} |x|^{-bp}f\eta^2(u^n)^{q-2}u + 4 \int_{\Omega} |x|^{-2a}|\nabla\eta|^2u^2(u_n)^{q-2}. \end{aligned} \quad (4.5)$$

Furthermore, an explicit calculation gives

$$\begin{aligned} |\nabla((u^n)^{\frac{q}{2}-1}u\eta)|^2 &\leq \frac{(q+4)(q-2)}{4}(u^n)^{q-2}\eta^2|\nabla u^n|^2 + 2(u^n)^{q-2}|\nabla u|^2\eta^2 \\ &\quad + 2(u^n)^{q-2}u^2|\nabla\eta|^2 + \frac{q-2}{2}(u^n)^q|\nabla\eta|^2. \end{aligned} \quad (4.6)$$

Let $C(q) := \min \left\{ \frac{1}{4}, \frac{4}{q+4} \right\}$. From (4.5) and (4.6) we get

$$\begin{aligned} C(q) \int_{\Omega} \frac{|\nabla((u^n)^{\frac{q}{2}-1}u\eta)|^2}{|x|^{2a}} &\leq 2(2+C(q)) \int_{\Omega} \frac{(u^n)^{q-2}u^2|\nabla\eta|^2}{|x|^{2a}} + C(q) \frac{q-2}{2} \int_{\Omega} \frac{(u^n)^q|\nabla\eta|^2}{|x|^{2a}} \\ &\quad + \int_{\Omega} \frac{f(x)}{|x|^{bp}} \eta^2 (u^n)^{q-2}u + \int_{\Omega} \frac{V(x)}{|x|^{bp}} \eta^2 (u^n)^{q-2}u^2. \end{aligned} \quad (4.7)$$

Estimate (4.4) applied to $\eta(u^n)^{\frac{q}{2}-1}u$ gives

$$\int_{\Omega} \frac{V^+(\eta(u^n)^{\frac{q}{2}-1}u)^2}{|x|^{bp}} \leq \frac{C(q)}{2} \int_{\Omega} \frac{|\nabla(\eta(u^n)^{\frac{q}{2}-1}u)|^2}{|x|^{2a}} + \ell \int_{\Omega \cap B_{\ell}(0)} \frac{(u^n)^{q-2}u^2\eta^2}{|x|^{bp}}. \quad (4.8)$$

By Hölder's inequality and convexity we arrive at

$$\int_{\Omega} \frac{|f|\eta}{|x|^{\frac{bp}{q}}} \frac{(u^n)^{q-2}u\eta}{|x|^{\frac{bp(q-1)}{q}}} \leq \frac{q-1}{q} \int_{\Omega} |x|^{-bp} \eta^{\frac{q}{q-1}} |u^n|^{q\frac{q-2}{q-1}} |u|^{\frac{q}{q-1}} + \frac{1}{q} \int_{\Omega} |x|^{-bp} |f|^q \eta^q. \quad (4.9)$$

We use (4.8) and (4.9) to estimate the terms with f and V in (4.7), then (1.3) yields

$$\begin{aligned} &\left(\int_{\Omega} |x|^{-bp} |u^n|^{(\frac{q}{2}-1)p} |u|^{p\eta^p} \right)^{\frac{2}{p}} \\ &\leq \frac{2\mathcal{C}_{a,b,N}(q-1)}{C(q)q} \int_{\Omega} |x|^{-bp} \eta^{\frac{q}{q-1}} |u^n|^{q\frac{q-2}{q-1}} |u|^{\frac{q}{q-1}} + \frac{2\mathcal{C}_{a,b}}{C(q)q} \int_{\Omega} |x|^{-bp} |f(x)|^q \eta^q \\ &\quad + \frac{2\ell\mathcal{C}_{a,b,N}}{C(q)} \int_{\Omega \cap B_{\ell}(0)} |x|^{-bp} \eta^2 |u^n|^{q-2} u^2 + \frac{4\mathcal{C}_{a,b}(2+C(q))}{C(q)} \int_{\Omega} |x|^{-2a} |u^n|^{q-2} u^2 |\nabla\eta|^2 \\ &\quad + \mathcal{C}_{a,b,N}(q-2) \int_{\Omega} |x|^{-2a} |u^n|^q |\nabla\eta|^2. \end{aligned}$$

Letting $n \rightarrow \infty$ in the above inequality (4.3) follows. Observe that if $u \in \mathcal{D}_a^{1,2}(\Omega)$ then we need not to use the cut-off η and the same analysis as above gives the estimate (4.3) for $\Omega' = \Omega$. The lemma is thereby proved. \square

Remark 4.2. By Vitali's theorem V belongs to $L^{p/(p-2)}(\Omega, |x|^{-bp})$ if and only if there exists ℓ such that (4.2) is satisfied. But the constant in (4.3) depends uniformly on ℓ and not on the norm of V in $L^{p/(p-2)}(\Omega, |x|^{-bp})$.

Proof of Theorem 1.2. We apply Lemma 4.1 with $f = 0$ and $V(x) = K(x)|u|^{p-2}$.

Starting with $q = p$, the lemma gives $u \in L_{\text{loc}}^{\frac{p^2}{2}}(\Omega, |x|^{-bp})$. Taking $q = \frac{p^2}{2}$, we find $u \in L_{\text{loc}}^{\frac{p^3}{4}}(\Omega, |x|^{-bp})$. Iterating the process, we obtain that $u \in L_{\text{loc}}^{p^{k+1}/2^k}(\Omega, |x|^{-bp})$ for any k . Let $k_0 \in \mathbb{N}$ be such that $(p/2)^{k_0} \geq 2(p-1)/(p-2)$, then after k_0 steps we find that $u \in L_{\text{loc}}^{\frac{2p(p-1)}{p-2}}(\Omega)$. Having this high integrability we may use Theorem 1.1 with $f(x) = K(x)|u|^{p-2}u$ to get the desired regularity of u . \square

APPENDIX A.

Lemma A.1. Let a, b and p satisfy (1.2) and $\varepsilon > 0$. Then we have

$$\left(\int_{B_{\rho}(x_0)} |x|^{-bp} \right)^{2/p+\varepsilon} \leq C_{(A.1)}(N) \rho^{-2+\varepsilon N} (\max(\rho, |x_0|))^{-\varepsilon bp} \int_{B_{\rho}(x_0)} |x|^{-2a}. \quad (A.1)$$

Proof. Let us distinguish two cases.

Case 1: $\rho \geq |x_0|/2$. Since $(N - bp)(2/p + \varepsilon) = N - 2 - 2a + \varepsilon(N - bp)$ we obtain

$$\left(\int_{B_\rho(x_0)} |x|^{-bp} \right)^{2/p+\varepsilon} \leq \left(\int_{B_{3\rho}(0)} |x|^{-bp} \right)^{2/p+\varepsilon} = C_1(N) \rho^{N-2-2a+\varepsilon(N-bp)}.$$

From the doubling property (2.1) and the fact that $B_\rho(0) \subset B_{4\rho}(x_0)$ we infer,

$$\begin{aligned} \rho^{\varepsilon(N-bp)-2} \int_{B_\rho(x_0)} |x|^{-2a} &\geq c \rho^{\varepsilon(N-bp)-2} \int_{B_{4\rho}(x_0)} |x|^{-2a} \\ &\geq c \rho^{\varepsilon(N-bp)-2} \int_{B_\rho(0)} |x|^{-2a} = C_2(N) \rho^{N-2-2a+\varepsilon(N-bp)} \end{aligned}$$

and the claim follows in Case 1.

Case 2: $\rho < |x_0|/2$. We have for all $x \in B_r(x_0)$ that $1/2 \leq |x|/|x_0| \leq 3$. Consequently,

$$\begin{aligned} \left(\int_{B_\rho(x_0)} |x|^{-bp} \right)^{2/p+\varepsilon} &\leq C_3(N) \rho^{N(2/p+\varepsilon)} |x_0|^{-2b-\varepsilon bp} \\ &\leq C_3(N) \rho^{N-2} |x_0|^{-2a} \rho^{2N/p-N+2} |x_0|^{-2(b-a)} \rho^{N\varepsilon} |x_0|^{-\varepsilon bp} \end{aligned}$$

From $r < |x_0|/2$ we get

$$\begin{aligned} \left(\int_{B_\rho(x_0)} |x|^{-bp} \right)^{2/p+\varepsilon} &\leq C_3(N) \rho^{N-2} |x_0|^{-2a} \rho^{N\varepsilon} |x_0|^{-\varepsilon bp} \\ &\leq C_4(N) \left(\int_{B_\rho(x_0)} |x|^{-2a} \right) \rho^{N\varepsilon} |x_0|^{-\varepsilon bp}, \end{aligned}$$

which ends the proof. \square

Lemma A.2. Suppose Φ be a nonnegative and nondecreasing functions on $[0, R]$ such that

$$\Phi(\rho) \leq A_1 \mu_a(B_\rho(x)) \mu_a(B_r(x))^{-1} \left(\frac{\rho}{r} \right)^{-\alpha} \Phi(r) + A_2 \mu_a(B_r(x)) r^{-\beta}, \quad (\text{A.2})$$

for any $0 < \rho \leq r \leq R$, where A_1, A_2, α and β are positive constants satisfying $\alpha < \beta$. Then for any $\gamma \in (\alpha, \beta)$ there exists a constant $C_{(A, \beta)} = C_{(A, \beta)}(A_1, \alpha, \beta, \gamma)$ independent of x and r such that for $0 < \rho \leq r \leq R$

$$\Phi(\rho) \leq C_{(A, \beta)} \left(\mu_a(B_\rho(x)) \mu_a(B_r(x))^{-1} \left(\frac{\rho}{r} \right)^{-\gamma} \Phi(r) + A_2 \mu_a(B_\rho(x)) \rho^{-\beta} \right). \quad (\text{A.3})$$

Proof. Fix $\gamma \in (\alpha, \beta)$ and set $\tau := \min(A_1^{-1/(\gamma-\alpha)}, 1/2)$. Then we have for $0 < r \leq R$

$$\Phi(\tau r) \leq \mu_a(B_{\tau r}(x)) \mu_a(B_r(x))^{-1} \tau^{-\gamma} \Phi(r) + A_2 r^{-\beta} \mu_a(B_r(x)).$$

Hence we may estimate for $k \in \mathbb{N}$

$$\begin{aligned} \Phi(\tau^{k+1} r) &\leq \mu_a(B_{\tau^{k+1} r}(x)) \mu_a(B_{\tau^k r}(x))^{-1} \tau^{\beta-\gamma} \\ &\leq \mu_a(B_{\tau^{k+1} r}(x)) \mu_a(B_r(x))^{-1} \tau^{-(k+1)\gamma} \Phi(r) + A_2 (\tau^k r)^{-\beta} \mu_a(B_{\tau^k r}(x)) \\ &\quad \cdot \underbrace{\sum_{j=0}^k \mu_a(B_{\tau^{k+1} r}(x)) \mu_a(B_{\tau^k r}(x))^{-1}}_{\leq 1} \underbrace{\mu_a(B_{\tau^{k-j} r}(x)) \mu_a(B_{\tau^{k-j+1} r}(x))^{-1}}_{\leq C_{(2.1)}(\tau) \text{ by (2.1)}} \tau^{(\beta-\gamma)j} \\ &\leq C_{(2.1)}(\tau) \mu_a(B_{\tau^{k+2} r}(x)) \mu_a(B_r(x))^{-1} \tau^{-(k+1)\gamma} \Phi(r) + \frac{A_2 C_{(2.1)}(\tau)}{1 - \tau^{\beta-\gamma}} (\tau^k r)^{-\beta} \mu_a(B_{\tau^k r}(x)) \end{aligned}$$

For $0 < \rho \leq r$ we may choose $k \in \mathbb{N}$ such that $\tau^{k+2}r < \rho < \tau^{k+1}r$ and obtain

$$\begin{aligned} \Phi(\rho) &\leq \Phi(\tau^{k+1}r) \\ &\leq C_{(2.1)}(\tau)\mu_a(B_\rho(x))\mu_a(B_r(x))^{-1}\left(\frac{\rho}{r}\right)^{-\gamma}\Phi(r) + \frac{A_2C_{(2.1)}^3(\tau)}{\tau(1-\tau^{\beta-\gamma})}\mu_a(B_\rho(x))\rho^{-\beta}. \end{aligned}$$

□

REFERENCES

- [1] H. Brézis and T. Kato. *Remarks on the Schrödinger operator with singular complex potentials*. J. Math. Pures Appl. (9), **58** (1979), no. 2, 137–151.
- [2] L. Caffarelli, R. Kohn and L. Nirenberg. *First order interpolation inequalities with weights*. Compositio Math., **53** (1984), no. 3, 259–275.
- [3] F. Catrina and Z.-Q. Wang. *On the Caffarelli-Kohn-Nirenberg inequalities: sharp constants, existence (and nonexistence), and symmetry of extremal functions*. Comm. Pure Appl. Math., **54** (2001), no. 2, 229–258.
- [4] E. Colorado and I. Peral. *Eigenvalues and bifurcation for elliptic equations with mixed Dirichlet-Neumann boundary conditions related to Caffarelli-Kohn-Nirenberg inequalities*. (2003). Preprint.
- [5] V. De Cicco and M. A. Vivaldi. *Harnack inequalities for Fuchsian type weighted elliptic equations*. Comm. Partial Differential Equations, **21** (1996), no. 9-10, 1321–1347.
- [6] E. B. Fabes, C. E. Kenig and R. P. Serapioni. *The local regularity of solutions of degenerate elliptic equations*. Comm. Partial Differential Equations, **7** (1982), no. 1, 77–116.
- [7] V. Felli and M. Schneider. *Compactness and existence results for degenerate critical elliptic equations* (2003). Preprint.
- [8] J. Heinonen, T. Kilpeläinen and O. Martio. *Nonlinear potential theory of degenerate elliptic equations*. The Clarendon Press Oxford University Press, New York (1993).

SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI, S.I.S.S.A., VIA BEIRUT 2-4, 34014 TRIESTE, ITALY

E-mail address: felli@sissa.it

MATHEMATISCHES INSTITUT, IM NEUENHEIMER FELD 288, D-69120 HEIDELBERG, GERMANY

E-mail address: mschneid@mathi.uni-heidelberg.de